



Royal College of Surgeons in Ireland  
**e-publications@RCSI**

Anatomy Articles

Department of Anatomy

1-1-2008

# Biomechanical comparison of the pullout properties of external skeletal fixation pins in the tibiae of intact and ovariectomised ewes.

Ben J. Keeley  
*University College Dublin*

Behnam Heidari  
*University College Dublin*

Nicholas J. Mahony  
*Trinity College Dublin*

Susan M. Rackard  
*University College Dublin*

Fergal J. O'Brien  
*Royal College of Surgeons in Ireland, [fjobrien@rcsi.ie](mailto:fjobrien@rcsi.ie)*

*See next page for additional authors*

## Citation

Keeley BJ, Heidari B, Mahony NJ, Rackard SM, O'Brien FJ, Lee TC, Brennan O, Kennedy OD, FitzPatrick D. Biomechanical comparison of the pullout properties of external skeletal fixation pins in the tibiae of intact and ovariectomised ewes. *Veterinary and Comparative Orthopaedics and Traumatology*. 2008;21(5):418-26.

This Article is brought to you for free and open access by the Department of Anatomy at e-publications@RCSI. It has been accepted for inclusion in Anatomy Articles by an authorized administrator of e-publications@RCSI. For more information, please contact [epubs@rcsi.ie](mailto:epubs@rcsi.ie).



---

**Authors**

Ben J. Keeley, Behnam Heidari, Nicholas J. Mahony, Susan M. Rackard, Fergal J. O'Brien, T Clive Lee, Orlaith Brennan, Oran D. Kennedy, and David FitzPatrick

---

## — Use Licence —

---

### **Attribution-Non-Commercial-ShareAlike 1.0**

#### **You are free:**

- to copy, distribute, display, and perform the work.
- to make derivative works.

#### **Under the following conditions:**

- Attribution — You must give the original author credit.
- Non-Commercial — You may not use this work for commercial purposes.
- Share Alike — If you alter, transform, or build upon this work, you may distribute the resulting work only under a licence identical to this one.

For any reuse or distribution, you must make clear to others the licence terms of this work. Any of these conditions can be waived if you get permission from the author.

Your fair use and other rights are in no way affected by the above.

---

This work is licenced under the Creative Commons Attribution-Non-Commercial-ShareAlike License. To view a copy of this licence, visit:

#### **URL (human-readable summary):**

- <http://creativecommons.org/licenses/by-nc-sa/1.0/>

#### **URL (legal code):**

- <http://creativecommons.org/worldwide/uk/translated-license>
-

# Biomechanical comparison of the pullout properties of external skeletal fixation pins in the tibiae of intact and ovariectomised ewes

B. J. Keeley<sup>1</sup>; B. Heidari<sup>2</sup>; N. J. Mahony<sup>3</sup>; S. M. Rackard<sup>1</sup>; F. J. O'Brien<sup>4</sup>; T. C. Lee<sup>3,4</sup>,  
O. Brennan<sup>4</sup>; O. D. Kennedy<sup>4</sup>; D. FitzPatrick<sup>2</sup>

<sup>1</sup>UCD School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin, Republic of Ireland;

<sup>2</sup>UCD School of Electrical, Electronic and Mechanical Engineering, Engineering and Material Sciences Centre, University College Dublin, Belfield, Dublin, Republic of Ireland;

<sup>3</sup>Trinity Centre for Bioengineering, Department of Mechanical Engineering, Trinity College, Dublin, Republic of Ireland;

<sup>4</sup>Department of Anatomy, Royal College of Surgeons in Ireland, Dublin, Republic of Ireland

## Summary

The pin-bone interface is the least stable component of the external skeletal fixator. Concerns exist regarding the ability to obtain adequate implant purchase in poor quality bone. Consequently, reduced bone quality has been viewed as a contra-indication for the use of external skeletal fixators. The aim of this study was to investigate the holding power of two different fixator pin designs in bone from entire and ovariectomised sheep. Thirty-two aged ewes were divided into two groups. Group 1 were controls, and Group 2 were ovariectomised (OVX). The ewes were sacrificed 12 months post-ovariectomy and five pairs of tibiae were harvested from each group. The holding power of cortical and cancellous fixator pins was assessed at five standardised locations on each tibia. An increase in mean cortical thickness was noted in the OVX group. The holding power of cancellous fixator pins was superior to that of cortical pins, irrespective of whether or not ovariectomy had been performed. Cancellous pins had an increased holding power in post ovariectomy bone compared to control bone. Cortical pin performance was not affected by ovariectomy. There was a lack of correlation between the incidence of insertional fractures of the far cortex and implant holding power. The results raise questions over the effectiveness of ovariectomy in establishing osteopaenic bone suitable for assessing implant performance, hence further investigations are warranted.

## Keywords

Implant, holding power, mechanical properties, cortical, cancellous

Vet Comp Orthop Traumatol 2008; 21: 418–426

Prepublished online April 11, 2008  
doi:10.3415/VCOT-07-09-0089

## Introduction

The pin-bone interface is the least stable component of the external skeletal fixator (1). As such, it is most often the determining factor in fixator longevity and subsequent clinical success or failure (2). Resistance to acute axial extraction (pullout strength) is an indirect measurement of integrity of the pin-bone interface, and is often referred to as 'holding power' (3). Holding power is known to be influenced by the method of pin insertion (4), pin design (5) and bone density (6, 7).

Low mineral density and deteriorated micro-architecture of bone are not only major risk factors for fractures, they also contribute to orthopaedic complications in fracture stabilisation. The attainment of adequate implant purchase remains a challenge in poor quality bone (8–10). Historically, pin-fixation has been contra-indicated in the stabilisation of such pathological bone (6, 11–14) however, recent *in vivo* studies have demonstrated that reduced quality bone no longer needs to be viewed as a contra-indication for the use of external skeletal fixators (15, 16).

The aspects of major interest in orthopaedic surgery are bone fragility, efficacy of implant fixation and bone healing. Various osteopaenic animal models have been established and used to investigate implant performance (17–20). The sheep has been proposed as an effective model (21–25) be-

cause ovine bone remodels and heals similarly to human bone (26). Decreased bone mineral density (23) and a reduction in the mechanical properties (27) of ovine bone have been demonstrated post ovariectomy. Increased bone turnover (26, 28), and reduced bone mass (29, 30) have also been reported. Biomechanical markers of bone formation like bone-specific alkaline phosphatase have been reported to increase in sheep after ovariectomy, indicating an increase in bone turnover similar to the human postmenopausal condition (23).

The aim of this study was to investigate the effect of ovariectomy on the holding power of threaded implants using an ovine osteopaenia model. This paper describes an *in vitro* assessment of threaded external skeletal fixator pins in tibiae harvested from ewes one-year post-ovariectomy, and compares the findings to a control group of similarly managed entire ewes.

## Materials and methods

### Surgical procedure

Thirty-two aged ewes were included in this project. Sixteen underwent ovariectomy (treatment group) while the remainder were maintained under similar conditions but did not have their ovaries removed (control group). Under Irish Government license, ovariectomy was performed through a ven-

Received September 17, 2007  
Accepted February 4, 2008

tral midline laparotomy under general anaesthesia, induced with 5% sodium thiopentone<sup>a</sup> (5–20 mg/kg to effect i.v.), intubated and maintained on isoflurane<sup>b</sup> (1.5–3%) in the surgical facilities at the UCD Lyons research farm. An analgesic (flunixin meglumine<sup>c</sup>, 2.2 mg/kg i.v.) was administered perioperatively and again 24 hours post-operatively by intravenous injection. All of the procedures were performed by trained veterinary surgeons. One-year post ovariectomy, all of the sheep were euthanatized for scientific reasons and the skeletons were harvested.

### Assessment of bone quality

A wide variety of histomorphological and biomechanical tests were carried out on cortical and cancellous bone samples collected from both the control and post-ovariectomy groups. Micro-computed tomography imaging was performed to generate data relating to the histomorphology and trabecular microarchitecture of the specimens. Static compression testing was carried out in order to obtain information about the specimens' mechanical properties and epifluorescence microscopy was utilized to quantify the degree of bone turnover. The results demonstrated that the sheep in the ovariectomy group (OVX) had changes in their cortical and cancellous bone quality consistent with the onset of osteoporosis (28, 30, 31). While there was a non-significant reduction in cancellous bone mineral density in the OVX group, there was a significant reduction in the modulus and ultimate compressive strength of the OVX cancellous bone (30). Cortical bone turnover and cortical porosity were significantly increased in the OVX bone although this did not translate into a significant reduction in compressive bone strength (28).

**Table 1**  
Specifications of each external fixator pin design

	Cortical pin	Cancellous pin
Material	316 L stainless steel	316 L stainless steel
Minor (shank) diameter (mm)	3.2	3.2
Major (thread) diameter (mm)	4.0	4.8
Thread pitch (mm)	0.9	1.45
Recommended drill bit (mm)	3.1	3.1
Thread length (mm)	35	35
Overall length (mm)	130	130

### Bone preparation

The left and right tibiae were randomly collected from five ewes in the control group (CON) and five ewes from the group OVX. Within eight hours of euthanasia, the bones were stripped of their soft tissue attachments and dual energy X-ray absorptiometry<sup>d</sup> (DEXA) was used to determine the bone mineral density (BMD) at specific points along each tibia. The bones were individually wrapped in saline (0.9% NaCl) soaked towels, packaged in double plastic bags, and stored at –20°C. The specimens were numbered and the investigators performing the biomechanical testing were blinded to the group (CON or OVX) from which each specimen originated. Five pairs of tibiae were thawed at room temperature 24 hours before pin insertion and mechanical pullout testing. All of the specimens were kept moist with saline at all times. Several thawing and refreezing sequences do not change the mechanical properties of bone (32).

### Pin selection

Commercially available stainless steel cortical and cancellous, bicortical, end-threaded, positive profile pins (Veterinary Instrumentation, Sheffield, UK) were inserted into one of five locations on each tibia. All of the pins were new and were used only once during the study (Table 1).

The toss of a coin was used to randomly select which pin design (cortical or cancellous) was inserted at each of the five desig-

nated sites on the left tibia of each pair of tibiae. The alternate pin type was then placed at the equivalent site on the contralateral (right) tibia.

### Pin insertion

Tibial positioning, and pin insertion were performed by the same investigator (B. Keeley). Each tibia was mounted in an identical position onto a custom-designed frame. Plastic cable ties and linseed oil putty (Siroflex Ltd Valliance Works, Leeds, UK) were used in order to align each tibia so that its sagittal plane was parallel to the frame. The fixture was then turned through 90 degrees and held in a vice to allow insertion of the fixator pins in a medial-to-lateral direction. Five pin insertion sites were selected at 5%, 25%, 50%, 75%, and 95%, respectively, of the distal to proximal length of the tibia (Fig. 1).

Prior to insertion of each pin a 3.1-mm pilot hole<sup>e</sup> was drilled perpendicular to the longitudinal axis of the tibia in a medial-to-lateral direction using a high-speed (600 rpm) bench mounted drill (Model # HDP600B, SIP [Industrial Products] Ltd., Loughborough, UK). A new drill bit was used for every eighth hole drilled. All of the pins were placed by the same investigator (B. Keeley), in accordance with the manufacturer's guidelines. Pin placement was performed using a slow-speed (<150 rpm) power drill (Hitachi Impact Drill DV18V, Hitachi Europe Ltd., Maidenhead, UK) and insertion was stopped once the entire tro-


<sup>a</sup> Thiovet<sup>™</sup>, Novartis Animal Health Ltd., Brintree, UK.

<sup>b</sup> Forane<sup>™</sup>, Abbott Laboratories Ltd., Maidenhead, UK.

<sup>c</sup> Flunixin<sup>™</sup> injection, Norbrook Laboratories Ltd., Newry, Northern Ireland.

<sup>d</sup> Hologic QDR-4500<sup>™</sup> Elite, Hologic, Bedford, MA, USA.

<sup>e</sup> Sticktite<sup>™</sup> Drill Bit, IMEX<sup>™</sup> Veterinary, Inc., Longview, TX, USA.



**Fig. 1**  
Craniocaudal radiograph of a right tibia post bio-mechanical testing and implant removal. A radio-opaque marker was placed to allow quantification of any radiographic magnification.

char tip of the pin had penetrated the far cortex. A second investigator observed pin placement and noted when the correct pin insertion depth was achieved, allowing one continuous insertion process.

## Mechanical testing

A stainless steel flat washer (R.A. Poole & Co. Ltd., Chessington, UK) (internal diameter 18-mm, external diameter 35 mm) was affixed to the medial surface of the bone at each pin insertion point using methyl methacrylate (Demotec 95, Demotec Siegfried Demel, Nidderau, Germany). Washers were placed at 90° to the long axis of the pin in order to ensure a planar bearing surface for pullout testing. A spirit level was used to guarantee that the washers were level and that care was taken to exclude methyl methacrylate from within the internal diameter

of each washer. Biomechanical testing was performed on the same day as pin insertion. The test specimen was positioned in a test fixture from which a counter force was produced against the washer. Pins were axially extracted with a constant extraction speed of 5 mm/minute, as defined by The American Society for Testing and Materials (ASTM) guidelines, using a universal testing machine (Hounsfield THE 050KS Test Equipment, Salfords, UK). The tibia was repositioned for each pullout test to accomplish exact alignment of each pin at all five tibial locations. The results of all pin pullout tests were recorded on a personal computer, and the ultimate load required to cause specimen failure was determined from the load displacement curves recorded (QMat Professional, Tinius Olsen Ltd., Salfords, UK); the ultimate load being marked by a clear drop off in the curve.

## Radiography

After testing, the pins were removed using a slow speed (<150 rpm) electric drill (Siroflex Ltd. Valliance Works, Leeds, UK). A craniocaudal radiographic projection of each tibia was performed using digital radiography (Kodak CR500, Kodak Limited, Hemel Hempstead, UK). A radio-opaque marker was placed next to each tibia to allow quantification of any radiographic magnification. At each diaphyseal insertion site the cortical bone thickness and bone diameter were measured using the Kodak CR500 software. Cortical thickness was defined as the sum total of the width of the near and far cortices in the central cross-section of each diaphyseal insertion site. Pin tract length alone was measured at the metaphyseal insertion sites, as accurate assessment of the cortical bone thickness at these locations was deemed impossible.

## Holding power

The holding power was determined for each pin at each of the insertion sites (33). For the purposes of this article, holding power was defined as yield strength per millimetre of cortical bone thickness at the diaphyseal insertion sites or yield strength per millimetre of pin tract at the metaphyseal insertion sites.

## Mode of failure

Mode of specimen failure was categorised using visual examination of the near and far cortices of each tibia during and after pullout testing and from post-testing radiographs. Mode of specimen failure was categorised as: pin slippage within the universal testing machine, bone failure, implant failure, or failure at the bone-implant interface (shearing of bone around the pin threads). Failure was defined as an acute reduction in the ability of the pin to resist axial extraction from the bone specimen, visualised as a clear drop off in the load displacement curve.

## Statistical analysis


Pullout data were expressed as mean  $\pm$  SD. Student's *t*-test assuming equal sample variance was used to compare pin pullout data.  $P < 0.05$  was considered statistically significant.

## Results

### DEXA analysis

A significant difference in BMD was not noted between groups at any of the pin insertion sites. The mean global BMD was  $0.93 \pm 0.09$  g/cm<sup>2</sup> and  $1.00 \pm 0.04$  g/cm<sup>2</sup> for the CON and OVX groups, respectively.


**Fig. 2**  
Cortical thickness data. A graph comparing the mean cortical thickness and standard deviation for each of the three diaphyseal pin insertion sites. Site 2  $p = 0.029$ , site 3  $p = 0.042$ , and site 4  $p = 0.002$ , respectively.



### Cortical thickness


Evaluation of the cortical bone thickness at the diaphyseal pin insertion sites revealed a large variation in morphology between the different bone specimens and at different implant insertion sites. The maximum cortical thickness in the diaphyseal segments

varied between 4.4 mm and 8.4 mm in the CON group, and 5.4 mm and 8.8 mm in the OVX group. The mean cortical thickness was significantly greater in the OVX group at all three diaphyseal insertion sites. A significant difference in pin tract length (bone diameter) was not observed between the CON and OVX groups (Figs. 2 and 3).




**Fig. 3** Box and whisker plot illustrating absolute yield strength at the different pin insertion sites. A graph comparing the mean holding power and standard deviation of cortical and cancellous pins in CON and OVX bone






**Fig. 4** Graph comparing the mean holding power and standard deviation of cortical and cancellous pins in CON and OVX bone.



**Fig. 5** Photograph documenting the generation of a butterfly fragment on the medial aspect (near cortex) of an ovine tibia



**Fig. 6** Radiograph documenting the generation of a butterfly fragment on the medial aspect (near cortex) of an ovine tibia

## Holding power

According to the axial extraction forces recorded, cortical and cancellous pins placed in the CON group behaved differently from those in the OVX group. Implant holding power was significantly greater at site 5 ( $p=0.016$ ) in the OVX group compared to the CON group irrespective of pin design. The same strong trend was noted at site 2 ( $p=0.078$ ). In OVX bone, cancellous pins had a greater holding power than cortical pins at insertion sites 2 ( $p=0.048$ ), 3 ( $p=0.023$ ) and 5 ( $p=0.039$ ). An association between extraction force of cortical pins and bone quality was not observed.

## Pin design

The 3.2/4.8 mm cancellous pin was superior ( $p<0.05$ ) at insertion sites 3 and 5 compared to the 3.2/4.0 mm cortical pin, irrespective of bone quality. The same trend was also noted at insertion site 1 ( $p=0.071$ ). Cancellous pins had a statistically significant ( $p<0.05$ ) increased holding power in OVX bone at pin insertion site 5 ( $p=0.001$ ), and the same trend was noted at pin site 2 ( $p=0.077$ ) and site 4 ( $p=0.060$ ). There was not a statistically significant difference between the holding powers of cortical or cancellous fixator pins in CON bone (Fig. 4).

## Failure mode

In all pullout tests, bone failure occurred without implant breakage. Pin pullout testing of both pin types consistently resulted in bone shearing along a line parallel to the outer edge of the pin's thread. In all diaphyseal regions, there was shearing of the pin-bone interface at the far cortex with generation of a butterfly fragment at the near cortex (Figs. 5 and 6).

## Coincidental findings

An audible crack was heard during advancement of 33% of both cortical and cancellous pins through the far cortex. This sound corresponded to visually evident chip fractures



adjacent to the exit holes of the enhanced threaded pins and was only heard during implant placement at the diaphyseal insertion sites (Table 2).

## Discussion

For uneventful healing of long bone fractures treated by external fixation, certain conditions are required of the bone/implant construct, namely the preservation of the capacity of the bone to heal and of a mechanically sound fixation. Fatigue of the implant, implant loosening and bone resorption may jeopardise optimal maintenance of a stable fracture reduction. Implant loosening is the result of a complex process of both mechanical and biological events that are initiated by microstructural trauma which occurs during the insertion of threaded implants (34, 35). Static preloading and cyclic loading of the implant by the patient propagates micro-cracks causing further implant loosening (35). Biological alterations, owing to vascular and thermal insult during implant insertion, result in the death of peri-implant bone (34), which further compromises implant holding power and fracture stability.

This study compared pin thread variables (cortical and cancellous designs) at metaphyseal and diaphyseal locations of ovine tibiae. The tibia was selected because of the relative frequency of external skeletal fixator usage on this bone. The use of paired tibiae for comparative implant pullout testing was considered to be an appropriate comparison technique in order to assess biomechanical properties of the two pin designs.

All biomechanical tests ended in failure of the pin-bone interface; bone shearing as a cylinder with a diameter equal to the outer thread diameter of the corresponding pin. Occasionally, a small cone of bone fractured from the near cortex and remained attached to the pin thread. This was due to the shear strength of the portion of bone avulsed being less than or equal to the implant holding power at this site (3, 36). Implant holding power is dependent on the major diameter of the pin, as well as the pin's minor diameter, thread depth (defined as the dis-

**Table 2**  
Incidence of insertional fractures of the far cortex

		Site 2	Site 3	Site 4	Total
CON	Cortical	0	3	2	5
	Cancellous	1	4	2	7
OVX	Cortical	0	1	1	2
	Cancellous	0	6	1	6
	Total	1	13	6	20

tance from the apex to the base of the thread), extent of cortical bone engaged by the thread, density and shearing strength of the bone and the size of the pilot hole drilled (12, 13, 37, 38). It has been reported that the major diameter of a threaded implant needs to be considered as the most important implant-related factor influencing pullout force; other factors such as thread design, pitch, and minor diameter having been shown to have a lesser effect on pullout forces (5, 36, 39). The difference in major diameter between pin designs tested could explain the improved performance of cancellous pins compared to cortical pins.

Cancellous pins have been specifically designed for use in the metaphyseal and epiphyseal regions of long bones, where the cancellous bone has thin trabeculae (40). In this study, there was a trend towards cancellous pins having a greater holding power at the distal metaphyseal insertion sites, compared to cortical pins. However, the cancellous pins performed significantly better than cortical pins when placed at the proximal metaphyseal site of OVX bone. This improved holding power can most likely be explained by the design characteristics of cancellous pins. The larger thread depth and pitch, and a larger ratio between the major and minor pin diameters (37) allows a greater volume of bone to reside between threads and increases pin to bone surface contact (41).

There was not a statistically significant ( $p < 0.05$ ) difference between the maximum holding power of the pin designs at the metaphyseal insertion sites in control bone, despite the difference in the thread diameter. It has been reported that in regions with a cortical thicknesses of less than 1.5 mm, cancellous density determined the ultimate pullout load, while in regions with cortices thicker than 1.5 mm, cortical thickness

alone significantly influenced the holding capacity of an implant (42). At metaphyseal locations, Marti and Roe (1999) reported that there is only a small amount of thread-bone contact between the trabeculae and the pin, the primary site of engagement appearing to be the thin cortical shell (43). The influence of pitch and thread depth on the pullout strength of implants in thin cortical bone warrants further study.

Insertional trauma associated with the use of cancellous pins has been seen as a contraindication for their placement in diaphyseal bone (44) although the results of this *in vitro* study do not support this finding. The fracturing of the far cortex during implant insertion was noted in one third of all cases in this study, and was observed with both types of pin. This phenomenon has previously been reported following the use of enhanced threaded pins with and without the pre-drilling of a pilot hole prior to pin insertion (2, 33). A higher incidence of insertional fractures was noted when inserting cancellous pins at the mid diaphyseal insertion site, or when placing both cortical and cancellous pins in control bone. Although insertional torque was not measured in this study, it could be speculated that the greater major diameter of the cancellous pin resulted in increased insertional torque and a subsequent increased incidence of fractures of the far cortex. No correlation between the presence of mild to moderate fracturing of the far cortex and implant holding power was detected in this study.

Bone strength can be defined as the ability of a bone to endure the application of force without yielding or breaking. It 'reflects the integration of two main features: bone density and bone quality' (The National Institutes of Health Consensus Statement, 2000). With the advent of dual energy X-ray absorptiometry (DEXA)

scanning, the relationship between bone density and bone failure has been extensively investigated, including studies that prove a direct relationship between bone density and bone strength (45). A universally accepted definition of bone quality does not exist. Several factors may be involved; the most important is probably the micro-architecture of bone. It seems reasonable to view bone quality as a set of characteristics unrelated to bone mineral density that influence bone strength (46). The quality of bone is determined by many factors, including trabecular and cortical microarchitecture, bone turnover, the degree of mineralization of the bone matrix, and the amount of microdamage present (46).

Previous studies have shown that bone density has a major influence on implant holding strength, with only minor improvements being reached following modification of thread shape, pitch, and depth (6, 7). In this study, ovariectomy did not affect cortical pin holding power when compared with control bone. Additionally, cancellous pins had an increased holding power in OVX bone. The increase in cortical bone thickness noted in the OVX group is the most likely explanation for this improved performance. The relationship between cortical thickness and the surface area of the pin-bone interface has been previously reported (3, 11, 35, 36, 38, 39). Cortical bone thickness has a direct effect on the force required to extract a pin from the pin-bone specimen (47), implant holding increasing linearly and rapidly with increased thickness of cortex engaged by the thread (39).

The inability to detect a difference in bone mineral density (BMD) and altered implant holding power between the two study groups raises questions concerning the efficacy of this osteopaenia model. The significant increase in diaphyseal cortical thickness in the OVX group is one possible explanation for the inability of DEXA to differentiate between the two study groups. Bone remodelling secondary to reduced bone quality may have resulted in a compensatory increase in regional bone quantity. McNamara et al. (48) reported microtensile testing of individual trabeculae collected from the proximal tibia of control and ovariectomised rats. One of the group's main

findings was that the Young's modulus and yield strength of the cancellous bone from the ovariectomised group was higher than that from the control bone. This suggests that the biological system is compensating for the lower bone mass post ovariectomy by increasing the strength and stiffness of the bone that does remain. This may be seen as a 'Wolff's law' type adaptation where a homeostatic strain is maintained on the mechanosensitive cells by increasing the stiffness of the tissue when the amount of it has reduced. Adaptive bone remodelling seems to be a viable explanation for the increase in cortical bone thickness observed in this study, i.e. an increase in cortical bone thickness compensated for a decrease in cortical bone quality. The bone quantity value obtained using DEXA is based on a two-dimensional measurement of area and does not consider the volume of the bone. Consequently DEXA is unable to differentiate between thin cortices composed of dense bone and thicker cortices composed of low-density bone (49). It has been hypothesised that when using DEXA, quantitative changes in cancellous bone are being disguised by increases in cortical bone thickness (30).

Other possible explanations for the increase in cortical thickness should also be considered. Arens et al. (50) recently demonstrated that marked seasonal variations in BMD and biochemical markers of bone turnover occur in entire sheep, characterised by increasing bone mass in the summer and decreasing bone mass in the winter. Our study commenced and concluded during the summer months but it is conceivable that increased vitamin D production secondary to improved weather conditions prior to culling may have affected cortical thickness. It does, however, seem very unlikely that seasonal variations affected the OVX group and not the CON group. Campbell et al. (51) have previously reported significant differences in BMD in sheep. The increase detected in cortical thickness could potentially be explained by chance variation between populations.

It has been shown that short-term (one year) oestrogen depletion in the ewe can result in structurally significant changes in bone density within the cortical bone of the radial diaphysis (52). Kennedy et al. (28)

analysed intracortical bone turnover, intracortical porosity and level of resorption of bone harvested from the left metatarsal of the ewes included in this study. Cortical bone turnover was significantly increased in the OVX group at six, nine and 12 months. Increased intracortical porosity and resorption was also demonstrated in the OVX group 12 months post-ovariectomy ( $p < 0.05$ ). Bone quality parameters were significantly altered in the metatarsal cortical bone and it would seem fair to assume that this deterioration in bone quality is generalised. Despite the detection of significant alterations in bone turnover parameters and porosity in Kennedy's study, mechanical testing failed to recognise changes in the compressive strength of the ovine cortical bone 12 months post-ovariectomy. The findings in this study fail to support the theorem that OVX bone has a reduced strength. It may be more appropriate to view post-ovariectomy bone as simply different rather than of reduced strength.

It is also worth considering the argument that ovariectomised sheep may not be a good model for osteoporosis (20, 53). The requirements for an animal model simulating the behaviour of osteopaenia bone during fracture treatment are different from those for pharmacological testing (20). Most animal models simulating the osteoporotic condition have been used to test drugs that have been developed to treat osteoporosis. Bone fragility, efficacy of implant fixation and bone healing must all be taken into consideration in the field of orthopaedics and traumatology. It is generally accepted that bone fragility is reflected in decreased BMD and bone mechanical properties (20). We failed to demonstrate significant changes in BMD or ultimate compressive strength in the cortical bone of ovine tibiae 12 months post-ovariectomy.

This biomechanical study provides a comparison of the holding power, measured as pullout strength, of two commercially available fixation pin designs. Implant holding power is a relatively simple, reliable, repeatable *in vitro* test but it is not without its limitations as holding power is only one of the components that govern how pins maintain fracture stability clinically. More factors than biomechanical advantage deter-

mine the clinical success of a new fixation principle or a new implant. In particular, questions concerning the osseous response to pin implantation and loading must be addressed. An *in vivo* study would be necessary to determine how the differences in thermal and microstructural bone damage between the two pin types affect the long-term stability of the pin-bone interface.

Bone remodeling (resorption and formation) takes place throughout life and is orchestrated by a complex interplay of bone cells and factors that regulate the functions of the cells. The principle influences on the remodeling process are those derived from mechanical loading and hormonal signals. Caution must be expressed when extrapolating the results from this *in vitro* study to clinical cases of disuse osteopaenia. An absence of a continued load-bearing stress is accompanied by a marked cortical thinning secondary to decreased bone formation, whereas oestrogen depletion is associated with increased bone remodeling with resorption exceeding formation causing increased porosity. Further studies using *in vivo* models are required in order to investigate the effect that these differing bone responses have on implant holding power.

Postmenopausal bone remodelling is a well-established and heavily investigated phenomenon. Ovine animal models are widely accepted as an effective means of studying reduced bone quality (20). The significant effect that bone remodelling had on the holding power of the external fixator pins tested in this study highlights the need for further study of hormone-related bone remodelling. Continued investigations are necessary in order to test the efficiency and appropriateness of animal models for investigation of pin performance in osteoporotic bone.

#### Acknowledgements

This project was funded by the Higher Education Authority (HRB), under the Programme for Research in Third Level Institutions (PRTL) Cycle 3, and also by the Health Research Board of Ireland under Grant no. RP/2004/229. The authors thank Veterinary Instrumentation Ltd., Sheffield, UK, for donating the external skeletal fixator pins tested in this study. Thanks also go to Brian Cloak and the staff at the Lyons Research Facility, UCD for expert animal care and handling.

#### References

- Chao EY, Pope M. The mechanical basis of external fixation. In: Concepts in External Fixation. Seglison D, Pope M (eds). Orlando, FL: Grune & Stratton 1982; 13–39.
- McDonald DE, Palmar RH, Hulse DA et al. Holding power of threaded external skeletal fixation pins in the near and far cortices of cadaveric canine tibiae. *Vet Surg* 1994; 23: 488–493.
- Bechtol CD, Ferguson AB, Laing PG. Metals and Engineering in Bone and Joint Surgery. Baltimore: Williams & Wilkins 1959; 152–165.
- Egger EL, Hestand MB, Blass CE et al. Effect of fixation pin insertion on the bone-pin interface. *Vet Surg* 1986; 15: 246–252.
- Bennett RA, Egger EL, Hestand M et al. Comparison of the strength and holding power of 4 pin designs for use with half pin (type I) external skeletal fixation. *Vet Surg* 1987; 16: 207–211.
- DeCoster TA, Heetderks DB, Downey DJ et al. Optimising bone screw pullout force. *J Orthop Trauma* 1990; 4: 169–174.
- Halsey D, Fleming B, Pope MH et al. External fixator pin design. *Clin Orthop* 1992; 278: 305–312.
- Halverson TL, Kelley LA, Thomas KA et al. Effects of bone mineral density on pedicle screw fixation. *Spine* 1994; 19: 2415–2420.
- Wittenberg RH, Shea M, Swartz DE et al. Importance of bone mineral density in instrumented spine fusions. *Spine* 1991; 16: 647–652.
- McLain RF, McKinley TO, Yerby SA et al. The effect of bone quality on pedicle screw loading in axial instability. A synthetic model. *Spine* 1997; 22: 1454–1460.
- Ansell RH, Scales JT. A study of some factors which affect the strength of screws and their insertion and holding power in bone. *J Biomech* 1968; 1: 279–302.
- Numamaker DM, Perren SM. Force measurements in screw fixation. *J Biomech* 1976; 9: 669–675.
- Uhl RL. The biomechanics of screws. *Orthop Rev* 1989; 18: 1302–1307.
- Pennig D. The place of unilateral external skeletal fixation in the treatment of tibial fractures. *Int Orthop Trauma* 1991; 1: 161–176.
- Moroni A, Faldini C, Marchetti S et al. Improvement of the bone-pin interface strength in osteoporotic bone with use of hydroxyapatite-coated tapered external-fixation pins. *J Bone Joint Surg Am* 2001; 83: 717–721.
- Moroni A, Faldini C, Pegreff F et al. Osteoporotic pertrochanteric fractures can be successfully treated with external fixation. *J Bone Joint Surg Am* 2005; 87: 42–50.
- Mori H, Manabe M, Kurachi Y et al. Osseointegration of dental implants in rabbit bone with low mineral density. *J Oral Maxillofac Surg* 1997; 55: 351–361.
- Pan J, Shiota T, Ohno K et al. Effect of ovariectomy on bone remodelling adjacent to hydroxyapatite-coated implants in the tibia of mature rats. *J Oral Maxillofac Surg* 2000; 58: 877–882.
- Rocca M, Fini M, Giavaresi G et al. Tibial implants: biomechanical and histomorphometric studies of stainless steel and titanium screws hydroxyapatite-coated and uncoated in long-term ovariectomised sheep. *Int J Artif Organs* 2001; 24: 649–654.
- Egermann M, Goldhahn J, Schneider E. Animal models for treatment in osteoporosis. *Osteoporos Int* 2005; 16: S129–S138.
- Newman E, Turner AS, Wark JD. The potential of sheep for the study of osteoporosis: current status and comparison with other animal models. *Bone* 1995; 16: 277S–84S.
- Hornby SB, Ford SL, Mase CA et al. Skeletal changes in the ovariectomised ewe and subsequent response to treatment with 17 beta oestradiol. *Bone* 1995; 17(Suppl.): 389S–394S.
- Turner AS, Alvis M, Myers W et al. Changes in bone mineral density and bone-specific alkaline phosphate in ovariectomised ewes. *Bone* 1995; 17(Suppl.): 395S–402S.
- Turner AS, Mallinckrodt CH, Alvis MR, Bryant HU. Dose-response effects of estradiol implants on bone mineral density in ovariectomised ewes. *Bone* 1995; 17(4 Suppl): 421S–427S.
- Johnson RB, Gilbert JA, Cooper RC et al. Effect of estrogen deficiency on skeletal and alveolar bone density in sheep. *J Periodontol* 2002; 73: 383–391.
- Chavassieux P, Garnero P, Duboeuf F et al. Effects of a new selective oestrogen receptor modulator (MDL 103,323) on cancellous and cortical bone in ovariectomised ewes: a biomechanical, histomorphometric, and densitometric study. *J Bone Miner Res* 2001; 16: 89–96.
- Geusens P, Boonen S, Nijs J et al. Effect of salmon calcitonin on femoral bone quality in adult ovariectomised ewes. *Calcif Tissue Int* 1996; 59: 315–320.
- Kennedy OD. The effect of bone turnover on bone quality and material properties. [PhD thesis] Trinity College, Dublin, Ireland, 2007.
- Fini M, Pierini G, Giavaresi G et al. The ovariectomised sheep as a model for testing biomaterials and prosthetic devices in osteoporotic bone: a preliminary study on iliac crest biopsies. *Int J Artif Organs* 2000; 23: 275–281.
- Brennan O. The effect of osteoporosis on bone quantity, quality and strength. [PhD thesis]. Royal College of Surgeons, Dublin, Ireland 2007; Chapter 3, 61–87.
- Kelleher D, Kennedy OD, Rackard SM et al. Investigation into the effect of osteoporosis on trabecular bone: bone quality assessment of cored vertebral samples. Proceedings of the 12<sup>th</sup> Annual Conference of the Section of Bioengineering of the Royal Academy of Medicine in Ireland, (eds) McHugh P, O'Mahoney D, Fitzpatrick D. Galway, Ireland, 27–28 January, 2006, p80.
- Reed MR, Ali AM, Yang L et al. The biomechanical effects of freezing and thawing cancellous bone. *J Bone Joint Surg* 2000; 82B(Suppl. 2): 128.
- Anderson MA, Mann FA, Kinden DA et al. Evaluation of cortical bone damage and axial holding power of non-threaded and enhanced threaded pins placed with and without drilling of a pilot hole in femurs from canine cadavers. *J Am Vet Med Assoc* 1996; 208: 883–887.

34. Ling RSM. Observations on the fixation of implants to the bony skeleton. *Clin Orthop Rel Res* 1986; 210: 80–96.
35. Vangsness CT, Carter DR, Frankel VH. In vitro evaluation of the loosening characteristics of self tapped and nonself tapped cortical bone screws. *Clin Orthop Rel Res* 1981; 157: 279–286.
36. Schatzker J, Sanderson R, Murnaghan JP. The holding power of orthopaedic screws in vivo. *Clin Orthop* 1970; 108: 115–126.
37. Ferrara LA, Ryken TC. Screw pullout testing. In: *Mechanical Testing of Bone and the Bone-Implant Interface*. An YH, Draughn RA (eds). Boca Raton, FL: CRC Press 2000; 551–565.
38. Bynum D, Ledbetter WB, Boyd CL et al. Holding characteristics of fasteners in bone. *Expl Mech* 1970; 10: 474–480.
39. Koranyi E, Bowman CE, Knecht CD et al. Holding power of orthopaedic screws in bone. *Clin Orthop* 1972; 72: 283–286.
40. Schatzker J. Concepts of fracture stabilisation. In: *Bone in Clinical Orthopaedics*. Sumner-Smith G. Philadelphia: W.B. Saunders 1982; 387–398.
41. Johnson NL, Galuppo LD, Stover SM et al. An in vitro biomechanical comparison of the insertion variables and pullout mechanical properties of AO 6.5mm standard cancellous and 7.3mm self-tapping, cannulated bone screws in foal femoral bone. *Vet Surg* 2004; 33: 681–690.
42. Seebeck J, Goldhahn, Morlock MM et al. Mechanical behaviour of screws in normal and osteoporotic bone. *Osteoporos Int* 2005; 16: S107–S111.
43. Marti JM, Roe SC. An in vitro comparison of hollow ground and trocar points on threaded positive-profile external skeletal fixation pins in canine cadaveric bone. *Vet Surg* 1999; 28: 279–286.
44. Griffin H. IMEX™ Veterinary, Inc. 2006 product catalog p38 and personnel communication.
45. Bouxsein ML, Courtney AC, Hayes WC. Ultrasound and densitometry of the calcaneus correlate with the failure loads of cadaveric femurs. *Calcif Tissue Int* 1995; 56: 99–103.
46. Bouxsein G. Bone quality: where do we go from here? *Osteoporos Int* 2003; 14(Suppl. 5): 118–127.
47. Seebeck J, Goldhahn J, Stadel H et al. Effect of cortical thickness and cancellous bone density on the holding strength of internal fixator screws. *J Orthop Res* 2004; 22: 1237–1242.
48. McNamara LM, Prendergast PJ, Lyons CG et al. Strength of single rat trabeculae in normal, ovariectomised, and drug treated rat bone during aging. *Trans. 48<sup>th</sup> Orthopaedic Research Society meeting* 2003; 35.
49. Bouxsein ML. Determinants of skeletal fragility. *Best Pract Res Clin Rheumatol* 2005; 19: 897–911.
50. Arens D, Sigrist I, Alini M et al. Seasonal changes in bone metabolism in sheep. *Vet J* 2007; 174: 585–591.
51. Campbell AW, Bain WE, McRae AF et al. Bone density in sheep: genetic variation and quantitative trait loci localisation. *Bone* 2003; 33: 540–548.
52. Martin RB, Burr DB. A hypothetical mechanism for the stimulation of osteonal remodelling by fatigue damage. *J Biomech* 1982; 15: 137–139.
53. Sigrist IM, Gerhardt C, Alini M et al. The long-term effects of ovariectomy on bone metabolism in sheep. *J Bone Miner Metab* 2007; 25: 28–35.

#### Correspondence to:

Ben Keeley, BVM&S CertSAS MRCVS  
 Northwest Surgeons, Delamere House  
 Ashville Point, Sutton Weaver  
 Cheshire, WA7 3FW, UK  
 Phone: +44 1928 711 400, Fax: +44 1928 711 466  
 E-mail: bjk78@hotmail.com